

STRICT PARTITIONS OF MAXIMAL PROJECTIVE DEGREE

DAN BERNSTEIN

ABSTRACT. The projective degrees of strict partitions of n were computed for all $n \leq 100$ and the partitions with maximal projective degree were found for each n . It was observed that maximizing partitions for successive values of n “lie close to each other” in a certain sense. Conjecturing that this holds for larger values of n , the partitions of maximal degree were computed for all $n \leq 220$. The results are consistent with a recent conjecture on the limiting shape of the strict partition of maximal projective degree.

1. INTRODUCTION

Let $\lambda = (\lambda_1, \lambda_2, \dots)$ be a partition of n , denoted as usual by $\lambda \vdash n$. Let f^λ denote the number of standard tableaux of shape λ . f^λ is also the number of paths in the Young graph Y from the root (1) to λ , and it is also the degree of the irreducible character χ^λ of the symmetric group S_n .

A partition $\lambda = (\lambda_1, \dots, \lambda_r) \vdash n$ is said to be *strict* if $\lambda_1 > \lambda_2 > \dots > \lambda_r > 0$ for some r . In that case we write $\lambda \models n$. Let SY be the subgraph of the Young graph Y formed by the strict partitions. If $\lambda \models n$, let g^λ denote the number of paths in SY from the root (1) to λ . According to a theorem by Schur, g^λ is the degree of the projective representation of S_n corresponding to λ . It is also the number of standard young tableaux of *shifted shape* λ .

Vershik and Kerov [6, 7] have determined the asymptotic shape of the partition λ that maximizes f^λ as $|\lambda| = n \rightarrow \infty$. They have also shown that the same shape is also the asymptotic expected shape of a random partition with respect to the Plancherel measure. The latter result was reached independently by Logan and Shepp [4] as well. Through the Robinson-Schensted algorithm, the expected shape relates to the expected length of the longest increasing subsequence in a random permutation. For some recent developments related to this problem and the probability distributions involved, see [1, 2, 5].

More precisely, given the Young diagram of a partition $\lambda \vdash n$ where each box is 1×1 , shrink it along both axes by a factor of $\sqrt{n}$ to obtain the re-scaled diagram $\bar{\lambda}$ of total area 1. For each n , let $\lambda_{f_{\max}}^{(n)}$ be a partition $\lambda \vdash n$ with maximal f^λ , that is, $f^{\lambda_{f_{\max}}^{(n)}} = \max\{f^\nu \mid \nu \vdash n\}$. Through slight abuse of notation, where the maximizing partition is not unique for a given n , we shall take $\lambda_{f_{\max}}^{(n)}$ to read “any $\lambda \vdash n$ of maximal degree”.

Date: May 28, 2007.

Partially supported by the Israel Science Foundation (grant No. 947/04).

Theorem 1 ([6, 7]). *The limit shape as $n \rightarrow \infty$ of the re-scaled diagrams $\bar{\lambda}_{f_{\max}}^{(n)}$ exists, and is given by the two axes and by the parametric curve*

$$\begin{cases} x = \left(\frac{2}{\pi}\right) (\sin \Theta - \Theta \cos \Theta) + 2 \cos \Theta, \\ y = -\left(\frac{2}{\pi}\right) (\sin \Theta - \Theta \cos \Theta), \end{cases} \quad 0 \leq \Theta \leq \pi.$$




Figure 1

The Vershik-Kerov limit shape of Theorem 1 is shown in Figure 1.

The problem of determining the asymptotic shape of the partition λ which maximizes g^λ remains unsolved, and we are unaware of even partial characterizations of the shape. However, recently the following was conjectured.

Conjecture 2 ([3, Conjecture 8.2]). *The limit shape λ^* of the $\lambda \models n$ maximizing $2^{n-\ell(\lambda)} (g^\lambda)^2$ — and possibly maximizing g^λ — is given by the two axes and by the parametric curve*

$$\begin{cases} x = 2\sqrt{2} \cos \Theta, \\ y = \left(\frac{2\sqrt{2}}{\pi}\right) (\Theta \cos \Theta - \sin \Theta), \end{cases} \quad 0 \leq \Theta \leq \frac{\pi}{2}.$$




Figure 2

The conjectured limit shape of Conjecture 2 is shown in Figure 2. It was obtained from the Vershik-Kerov shape by bisecting it along the line $y = -x$, taking

the upper half, dilating it (so that its area become 1) and applying the shearing transformation $(x, y) \mapsto (x + y, y)$ (to bring the line $y = -x$ to the y axis).

In the next section, we give the results of computing the partition maximizing g^λ over all $\lambda \models n$ for $1 \leq n \leq 100$. We observe a property of successive maximizing partitions in the range $1 \leq n \leq 100$, and conjecture that it holds for all n . Assuming the conjecture, we compute the maximizing partitions for $100 < n \leq 220$. Our results are consistent with Conjecture 2 for both g^λ and $2^{-\ell(\lambda)} (g^\lambda)^2$.

2. RESULTS FOR $n \leq 100$

To compute g^λ , we used the following formula, due to Schur.

Theorem 3. *Let $\lambda = (\lambda_1, \dots, \lambda_r) \models n$. Then*

$$g^\lambda = \frac{n!}{\lambda_1! \cdots \lambda_r!} \prod_{1 \leq i < j \leq r} \frac{\lambda_i - \lambda_j}{\lambda_i + \lambda_j}$$

All computations were done with *Mathematica*.

For $1 \leq n \leq 100$, g^λ was computed for every $\lambda \models n$ and the partitions attaining the maximal value for each n were identified.

Observation 4 (Uniqueness of the maximum). For every $n \in [100] \setminus \{3, 11\}$, there exists a partition $\lambda = \lambda_{g \max}^{(n)} \models n$ such that $g^\lambda > g^\mu$ for all $\lambda \neq \mu \models n$.

Observation 5.

$$\lambda_{g \max}^{(100)} = (24, 20, 16, 13, 10, 8, 5, 3, 1).$$




Figure 3

Figure 3 shows the normalized diagram $\bar{\lambda}_{g \max}^{(100)}$ overlaid with the conjectured limit shape.

Recall that by Conjecture 2, the partitions maximizing g^λ are asymptotically equal to the partitions maximizing $2^{-\ell(\lambda)} (g^\lambda)^2$. For all strict partitions λ of $1 \leq n \leq 100$, $2^{-\ell(\lambda)} (g^\lambda)^2$ was computed and the partition maximizing it was denoted $\lambda_{2g \max}^{(n)}$. Comparing the results with the results for g^λ , the following was observed.

Observation 6. For all

$$n \in \{1, 2, \dots, 100\} \setminus \{3, 8, 16, 25, 26, 38, 51, 52, 53, 54, 69, 70, 88, 89, 90, 91\},$$

$$\lambda_{g \max}^{(n)} = \lambda_{2g \max}^{(n)}.$$

Definition 7 (d -successor). Let $\mu = (\mu_1, \dots, \mu_r) \models n$ and let $d \in \mathbb{N}$. The d -successors of μ are the elements of the set

$$N(\mu, d) := \{ \lambda = (\lambda_1, \dots, \lambda_s) \models n+1 \mid |\lambda_i - \mu_i| \leq d \quad 1 \leq i \leq s \}$$

We have observed the following.

Observation 8. For $1 \leq n < 100$, $\lambda_{g \max}^{(n+1)} \in N(\lambda_{g \max}^n, 1)$.

When $\lambda_{g \max}^{(n)}$ is not unique, read the above to mean “every $\lambda \models n+1$ of maximal projective degree is a 1-successor of every $\lambda \models n$ of maximal degree”.

3. A CONJECTURE AND RESULTS FOR $n \leq 220$

Based on the above observation, we conjecture the following.

Conjecture 9 (Maximizers are successors to a maximizers). *For all n , if $\lambda_{g \max}^{(n+1)} \in N(\lambda_{g \max}^n, 1)$.*

Assuming that the conjecture holds, $\lambda_{g \max}^{(n+1)}$ was computed for $100 \leq n < 220$ as follows: starting with $\lambda = \lambda_{g \max}^{(n)}$, for every $\mu \in N(\lambda, 1)$, the ratio $\frac{g^\mu}{g^\lambda}$ was computed and the μ maximizing the ratio was selected.

Observation 10. If Conjecture 9 holds for all $n < 220$, then

$$\lambda_{g \max}^{(220)} = (37, 32, 28, 24, 21, 18, 16, 13, 11, 8, 6, 4, 2).$$

Figure 4 shows the normalized diagram $\bar{\lambda}_{g \max}^{(220)}$ overlaid with the conjectured limit shape.

Figure 5 shows the ratio $\frac{(\lambda_{g \max}^{(n)})_1}{\ell(\lambda_{g \max}^{(n)})}$ for $1 \leq n \leq 220$, where the values above $n = 100$ are based on Conjecture 9. According to Conjecture 2, the ratio at the limit is π .




Figure 4




Figure 5

REFERENCES

- [1] J. Baik, P. Deift, and K. Johansson. On the distribution of the length of the longest increasing subsequence of random permutations. *J. Amer. Math. Soc.*, 12(4):1119–1178, 1999.
- [2] J. Baik, P. Deift, and K. Johansson. On the distribution of the length of the second row of a Young diagram under Plancherel measure. *Geom. Funct. Anal.*, 10(4):702–731, 2000.
- [3] D. Bernstein, A. Henke, and A. Regev. Maximal projective degrees for strict partitions. Preprint, 2006.
- [4] B. F. Logan and L. A. Shepp. A variational problem for random Young tableaux. *Advances in Math.*, 26(2):206–222, 1977.
- [5] R. P. Stanley. Recent progress in algebraic combinatorics. *Bull. Amer. Math. Soc. (N.S.)*, 40(1):55–68 (electronic), 2003. Mathematical challenges of the 21st century (Los Angeles, CA, 2000).
- [6] A. M. Vershik and S. V. Kerov. Asymptotics of the Plancherel measure of the symmetric group and the limit form of young tableaux. *Dokl. Akad. Nauk SSSR*, 233:1024–1027, 1977.
- [7] A. M. Vershik and S. V. Kerov. Asymptotic behavior of the maximum and generic dimensions of irreducible representations of the symmetric group. *Funktsional. Anal. i Prilozhen.*, 19(1):25–36, 96, 1985.

DEPARTMENT OF MATHEMATICS, THE WEIZMANN INSTITUTE OF SCIENCE, REHOVOT 76100, ISRAEL

E-mail address: dan.bernstein@weizmann.ac.il